





**CDMTCS  
Research  
Report  
Series**




**A new method for computing  
asymptotics of diagonal  
coefficients of multivariate  
generating functions**




**Alexander Raichev**  
Department of Computer Science,  
University of Auckland



**Mark C. Wilson**  
Department of Computer Science,  
University of Auckland



CDMTCS-296  
January 2007



Centre for Discrete Mathematics and  
Theoretical Computer Science

## Abstract

Let  $\sum_{\mathbf{n} \in \mathbb{N}^d} f_{\mathbf{n}} \mathbf{x}^{\mathbf{n}}$  be a multivariate generating function that converges in a neighborhood of the origin of  $\mathbb{C}^d$ . We present a new, multivariate method for computing the asymptotics of the diagonal coefficients  $f_{a_1 n, \dots, a_d n}$  and show its superiority over the standard, univariate diagonal method.

# A new method for computing asymptotics of diagonal coefficients of multivariate generating functions

ALEXANDER RAICHEV

Department of Computer Science,  
University of Auckland  
raichev@cs.auckland.ac.nz

MARK C. WILSON

Department of Computer Science,  
University of Auckland  
mcw@cs.auckland.ac.nz

## 1 Introduction

Let  $F(\mathbf{x}) = \sum_{\mathbf{n} \in \mathbb{N}^d} f_{\mathbf{n}} \mathbf{x}^{\mathbf{n}}$  be a complex power series that converges in a neighborhood of the origin but not on all of  $\mathbb{C}^d$ . Here  $\mathbf{x} = (x_1, \dots, x_d)$ ,  $\mathbf{n} = (n_1, \dots, n_d)$ , and  $f_{\mathbf{n}} \mathbf{x}^{\mathbf{n}} = f_{n_1, \dots, n_d} x_1^{n_1} \cdots x_d^{n_d}$ .

We wish to compute asymptotics for the diagonal coefficients  $f_{a_1 n, \dots, a_d n}$  for fixed positive integers  $a_1, \dots, a_d$ , a task often useful in enumerative combinatorics. For simplicity of presentation, we suppose that  $f_{\mathbf{n}} \geq 0$  and that  $F$  is rational, although much greater generality is possible. Thus far the only general method to extract diagonal asymptotics to be found in the literature is what we call the *standard diagonal method*. It consists of two steps: (1) find a closed form or defining equation for the diagonal generating function  $G(x) := \sum_{n \in \mathbb{N}} f_{a_1 n, \dots, a_d n} x^n$  via contour integration (in the style of [Fur67] and [HK71] and summarized in [Sta99, Section 6.3]); (2) apply univariate singularity analysis to  $G(x)$  (in the style of [FS] for instance) to compute the asymptotics. However, as elaborated below, this method is quite limited, working well only for main diagonals in two variables, that is, for computing the asymptotics of  $f_{nn}$ .

In this article we present a new, more versatile diagonal method. It consists of one step: apply multivariate singularity analysis (in the style of [PW02], [PW04], [BP]) directly to  $F(\mathbf{x})$  to compute the asymptotics.

## 2 Limitations of the Standard Diagonal Method

What is wrong with the standard diagonal method? It certainly works well for computing main diagonal asymptotics in two variables, as witnessed by the following example taken from [MZS04].

**Example 2.1 (*Zigzag-free Binary Words*).** The bivariate generating function

$$F(x, y) = \sum_{m, n} f_{mn} x^m y^n = \frac{1 + xy + x^2 y^2}{1 - x - y + xy - x^2 y^2}$$

counts the number of words over a binary alphabet,  $\{0, 1\}$  say, that have  $m$  zeros and  $n$  ones and do not contain zigzags, that is, the subwords 010 and 101. The main diagonal coefficients  $f_{nn}$ , then, count zigzag-free binary words with an equal number of zeros and ones.

To compute the asymptotics of  $f_{nn}$  using the standard diagonal method we proceed as follows. Since  $F(x, y)$  is rational and holomorphic in a neighborhood of the origin, for fixed  $x$  small enough  $F(x/t, t)$  will be rational and holomorphic as a function of  $t$  in some annulus about  $t = 0$ . Thus in that annulus it can be represented by a Laurent series whose constant term is  $[t^0]F(x/t, t) = \sum_{n \geq 0} f_{nn} x^n$ , the series we want. By Cauchy's Integral and Residue Theorems we have that for some circle  $\gamma_x$  about  $t = 0$

$$\begin{aligned} \sum_n f_{nn} x^n &= [t^0]F(x/t, t) \\ &= \frac{1}{2\pi i} \int_{\gamma_x} \frac{F(x/t, t)}{t} dt \\ &= \sum_k \text{Res}(F(x/t, t)/t; t = s_k), \end{aligned}$$

where the  $s_k$  are the “small” singularities of  $F(x/t, t)/t$ , that is, the ones satisfying  $\lim_{x \rightarrow 0} s_k(x) = 0$ . Since  $F$  is rational, these singularities are poles and algebraic functions of  $x$ , so that the residue sum, the diagonal generating function, is also an algebraic function of  $x$ .

In particular,

$$F(x/t, t)/t = \frac{1 + x + x^2}{-t^2 + (1 + x - x^2)t - x},$$

which has a single simple pole approaching zero as  $x$  approaches 0, namely,  $s = \frac{1}{2}(1 + x - x^2 - \sqrt{1 - 2x - x^2 - 2x^3 + x^4})$ . The residue, the diagonal generating function, is then

$$G(x) := \sqrt{\frac{x^2 + x + 1}{x^2 - 3x + 1}}.$$

Now, the singularity of  $G$  closest to the origin is  $\omega := (3 - \sqrt{5})/2$ , and its reciprocal is the exponential growth order of the coefficients of  $G$ . To determine the leading

subexponential factor we note that

$$G(x) \sim \left(1 - \frac{x}{\omega}\right)^{-1/2} \left( \sqrt{\frac{x^2 + x + 1}{-\omega \left(x - \frac{3+\sqrt{5}}{2}\right)}} \right)_{x=\omega} = \left(1 - \frac{x}{\omega}\right)^{-1/2} \frac{2}{\sqrt{\sqrt{5}}}$$

as  $x \rightarrow \omega$ , so that

$$f_{nn} = g_n \sim \omega^{-n} \frac{2}{\sqrt{\sqrt{5}}} \frac{n^{1/2-1}}{\Gamma(1/2)} = \left(\frac{2}{3 - \sqrt{5}}\right)^n \frac{2}{\sqrt{\sqrt{5}\pi n}}$$

by asymptotic transfer (see [FS, Chapter VI] for instance).  $\triangleleft$

However, the standard diagonal method encounters major problems off the main diagonal even in two variables, as illustrated in the next example adapted from [Sta99, Section 6.3].

**Example 2.2 (*Lattice Paths*).** The bivariate generating function

$$F(x, y) = \frac{1}{1 - x - y - xy},$$

whose coefficients  $f_{mn}$  are called the Delannoy numbers, counts the number of lattice paths from  $(0, 0)$  to  $(m, n)$  with allowable steps  $(1, 0)$ ,  $(0, 1)$ , and  $(1, 1)$ .

To compute the asymptotics of the general diagonal coefficients  $f_{an, bn}$  using the standard diagonal method, we fix  $x > 0$  small enough and try to find the small poles of

$$\begin{aligned} G(x) &:= \sum_n f_{an, bn} x^n \\ &= [t^0] F(x^{1/a}/t^b, t^a) \\ &= \frac{1}{2\pi i} \int_{\gamma_x} \frac{F(x^{1/a}/t^b, t^a)}{t} dt \\ &= \frac{1}{2\pi i} \int_{\gamma_x} \frac{t^{b-1}}{t^b - x^{1/a} - t^{a+b} - x^{1/a}t^a} dt \end{aligned}$$

Since  $t^b - x^{1/a} - t^{a+b} - x^{1/a}t^a$  has a zero of multiplicity  $b$  at  $t = 0$  when  $x = 0$ , it follows that  $F(x^{1/a}/t^b, t^a)/t$  has a single pole  $s$  of order  $b$  satisfying  $\lim_{x \rightarrow 0} s(x) = 0$ . Thus we have that

$$\begin{aligned} G(x) &= \text{Res}(F(x^{1/a}/t^b, t^a)/t; t = s) \\ &= \lim_{t \rightarrow c} \frac{1}{(b-1)!} D_t^{b-1} ((t-s)^b F(x^{1/a}/t^b, t^a)/t), \end{aligned}$$

where  $D_t$  is the derivative with respect to  $t$ . Patiently tracing through Leibniz's rule for the iterated derivative of a product, we could express this limit in terms of  $s$  and  $g(t) := (t-s)^{-b}(t^b - x^{1/a} - t^{a+b} - x^{1/a}t^a)$ , and use it to find an algebraic equation

satisfied by  $G$ . However, even with the help of Maple, this seems unlikely for general  $a$  and  $b$ .

Thus, already at the first step of the standard diagonal method, it seems we are thwarted, the cause of the problem being the parameters  $a$  and  $b$  occurring in the exponent: the larger they get, the greater the difficulty in finding a defining equation for  $G$ .  $\triangleleft$

So we should keep to main diagonal asymptotics when using the standard diagonal method. But even this poses problems in three or more variables, as illustrated by the next example.

**Example 2.3 (*Ternary Words*).** The trivariate generating function

$$F(x, y, z) = \frac{1}{1 - x - y - z}$$

counts the the number of words  $f_{lmn}$  over a ternary alphabet,  $\{0, 1, 2\}$  say, that have  $l$  zeros,  $m$  ones, and  $n$  twos (an easy consequence of the symbolic method as described in [FS]). An easy combinatorial argument shows that  $f_{lmn} = \binom{l+m+n}{l, m, n}$ , to which one could apply Stirling's formula and derive the asymptotics.

To compute the asymptotics of the main diagonal coefficients  $f_{nnn}$  using the standard diagonal method instead, we iterate the contour integration process. First

$$\begin{aligned} \sum_n f_{m,n,n} x^m y^n &= [t^0] F(x, y/t, t) \\ &= \frac{1}{2\pi i} \int_{\gamma_x} \frac{F(x, y/t, t)}{t} dt \\ &= \frac{1}{2\pi i} \int_{\gamma_x} \frac{1}{-t^2 + (1-x)t - y} dt, \end{aligned}$$

which has a single simple pole approaching zero as  $y$  and  $x$  approach 0, namely,  $s = \frac{1}{2}(1 + x - \sqrt{1 - 2x + x^2 - 4y})$ . The residue is then

$$G(x, y) := \frac{1}{\sqrt{1 - 2x + x^2 - 4y}}.$$

Second, since

$$\lim_{x \rightarrow 0} G(x/t, t) = \lim_{x \rightarrow 0} \frac{1}{\sqrt{1 - 2x/t + x^2/t^2 - 4t}} = \frac{1}{\sqrt{1 - 4t}},$$

$G(x/t, t)$  is a holomorphic function of  $t$  in some annulus about  $t = 0$ . Thus computing

$$\frac{1}{2\pi i} \int_{\gamma_x} \frac{G(x/t, t)}{t} dt = \frac{1}{2\pi i} \int_{\gamma_x} \frac{1}{t \sqrt{1 - 2x/t + x^2/t^2 - 4t}} dt$$

will give us the diagonal generating function. Proceeding, we employ Maple and find that two singularities of  $G(x/t, t)/t$  approach 0 as  $x$  approaches 0, namely

$$-\frac{1}{24}p(x) + \frac{q(x)}{p(x)} + \frac{1}{12} + \frac{1}{2}i\sqrt{3} \left( \frac{1}{12}p(x) + \frac{q(x)}{p(x)} \right)$$

and its conjugate, where  $p(x) = (-36x + 216x^2 + 1 + 24\sqrt{-3x^3 + 81x^4})^{1/3}$  and  $q(x) = x - \frac{1}{24}$ . While both of these singularities are algebraic in  $x$ , Maple can not compute the sum of their residues in reasonable time.

The problem is that the residue sum, that is, the diagonal generating function  $\sum \binom{3n}{n,n,n} x^n$ , is not algebraic (see [Sta99, Exercise solution 6.3] for instance). While it is D-finite—the diagonal of any rational function in any finite number of variables is D-finite [Lip88]—it is not trivial to find a closed form or defining differential equation for it. (In the present case we obtain, using the Maple package Mgfund, the defining differential equation  $27x(x-1)f''(x) + (54x-1)f'(x) + 6f(x) = 0$ , which has a hypergeometric solution). Moreover, even given the defining differential equation, we still require the theory of univariate singularity analysis of D-finite functions to compute the asymptotics. But this theory has not been worked out in general. Indeed, as Philippe Flajolet has informed us, certain aspects of it, such as the so-called connection problem, might not be computable!  $\triangleleft$

### 3 A New Diagonal Method

To transcend the limitations of the standard diagonal method, to go beyond main diagonal asymptotics of bivariate generating functions, we take a multivariate approach. When the diagonal method works, it does at least produce an explicit algebraic formula for the diagonal generating function, which our method here does not. However, it is normally much more useful to have an asymptotic expression for the coefficients. In order to achieve this, we can apply a multivariate singularity analysis directly to  $F(\mathbf{x})$ . Such an analysis has been recently developed by Baryshnikov, Pemantle, and Wilson in [PW02], [PW04], and [BP], the relevant parts of which we now summarize and adapt to our needs.<sup>1</sup>

Let  $\mathcal{D} \subset \mathbb{C}^d$  be the open domain of convergence of  $F$  and write  $F(\mathbf{x}) = I(\mathbf{x})/J(\mathbf{x})$  for some  $I$  and  $J$  holomorphic on an open domain  $\mathcal{D}'$  containing the closure  $\supsetneq \mathcal{D}$ , and relatively prime in the ring of holomorphic functions on  $\mathcal{D}'$ . In all examples in this paper, the representation of  $F$  as a quotient in fact holds on all of  $\mathbb{C}^d$ , but the extra generality is sometimes useful in applications. Let  $\mathcal{V}$  be the complex variety  $\{\mathbf{x} \in \mathbb{C}^d : J(\mathbf{x}) = 0\}$ .

A **critical point** of  $F$  for  $\mathbf{n} \in (\mathbb{N}^+)^d$  is a solution of

$$\begin{aligned} J(\mathbf{x}) &= 0 \\ n_d x_i J_i(\mathbf{x}) &= n_i x_d J_d(\mathbf{x}) \quad (i < d). \end{aligned}$$

Here  $n_i$  denotes component  $i$  of  $\mathbf{n}$ , and  $J_i$  denotes the partial derivative of  $J$  with respect to component  $i$  of the domain of  $J$ , conventions we adopt for all vectors and functions throughout and which should cause no confusion in context. Let  $\text{CRIT}(\mathbf{n})$  denote the set of all critical points of  $F$  for  $\mathbf{n}$ . For generic directions  $\mathbf{n}$ , this set is finite, being a

---

<sup>1</sup>As a whole, the multivariate singularity analysis developed in the aforementioned articles is much more general than what we present in this article, as it applies not only to rational functions but also to locally meromorphic functions. For a more complete summary including further examples, see the forthcoming article [PW].

zero-dimensional complex variety. The main situation in which  $\text{CRIT}(\mathbf{n})$  is infinite occurs when  $J$  defines a binomial variety  $\{\mathbf{x} \mid \mathbf{x}^{\mathbf{a}} - \mathbf{x}^{\mathbf{b}}\}$ , in which case  $\text{CRIT}(\mathbf{n})$  is empty for all but one direction and uncountable otherwise. Such examples can be analysed by a variant of the methods shown here.

A **contributing point** of  $F$  for  $\mathbf{n}$  is a critical point that influences the asymptotics of the coefficients of  $F$  in the direction of  $\mathbf{n}$ . Let  $\text{CONTRIB}(\mathbf{n})$  denote the set of all such points. While  $\text{CONTRIB}(\mathbf{n})$  is ill-defined here, its functional role will become clear from the next two theorems.

**Theorem 3.1.** If  $\text{CRIT}(\mathbf{n})$  is finite, then

- $\text{CRIT}(\mathbf{n})$  contains exactly one point, call it  $\mathbf{c}$ , that lies in the positive orthant of  $\mathbb{R}^d$ , and  $\mathbf{c} \in \text{CONTRIB}(\mathbf{n})$ ;
- all other members of  $\text{CONTRIB}(\mathbf{n})$  must lie on the same torus as  $\mathbf{c}$ ;
- in the case where  $J = 1 - P$  for some aperiodic power series with nonnegative coefficients  $P$ ,  $\text{CONTRIB}(\mathbf{n}) = \{\mathbf{c}\}$ .

Here, the **torus** of a point  $\mathbf{c} \in \mathbb{C}^d$  is the set  $\{\mathbf{x} \in \mathbb{C}^d : \forall i \leq d \ |x_i| = |c_i|\}$ , and a power series is **aperiodic** if the  $\mathbb{Z}$ -span of its monomial vectors is all of  $\mathbb{Z}^d$ .

A point  $\mathbf{c} \in \mathcal{V}$  is a **smooth point** if  $\mathcal{V}$  is a smooth complex manifold in a neighborhood of  $\mathbf{c}$ , or equivalently, if  $J_i(\mathbf{c}) \neq 0$  for some  $i$  (see [BK86, page 363] for instance). For simplicity of presentation, we deal only with smooth points in this article. This is the generic case, although interesting examples are not always generic. For more on the case of non-smooth points, see [PW04].

**Theorem 3.2.** Let  $\mathbf{n} = (a_1 n, \dots, a_d n)$  for some  $a_1, \dots, a_d \in \mathbb{N}^+$ . If  $\text{CONTRIB}(\mathbf{n})$  consists of a single smooth point  $\mathbf{c}$  such that  $c_d J_d(\mathbf{c}) \neq 0$  and  $c_d$  is a simple zero of  $x_d \mapsto J(c_1, \dots, c_{d-1}, x_d)$ , then

$$f_{\mathbf{n}} = \mathbf{c}^{-\mathbf{n}} \left( \sum_{j=0}^{j'} b_j n_d^{(1-d-j)/2} + O\left(n_d^{-d-j'}\right) \right)$$

for some  $b_j \in \mathbb{C}$  as  $n \rightarrow \infty$ . If  $\text{CONTRIB}(\mathbf{n})$  consists of a finite set of smooth points  $\mathbf{c}_1, \dots, \mathbf{c}_r$  each satisfying the hypotheses above, then

$$f_{\mathbf{n}} = \sum_i \mathbf{c}_i^{-\mathbf{n}} \left( \sum_{j=0}^{j'} b_{ij} n_d^{(1-d-j)/2} + O\left(n_d^{-d-j'}\right) \right)$$

as  $n \rightarrow \infty$ .

**Theorem 3.3.** In Theorem 3.2,

$$b_0 = \frac{I(\mathbf{c})}{-c_d J_d(\mathbf{c}) \sqrt{(2\pi)^{d-1} h(J, \mathbf{c})}},$$



assuming  $h(J, \mathbf{c}) \neq 0$ . Here  $h(J, \mathbf{c})$  is the determinant of the matrix  $\mathbf{H}$ , whose entries are

$$\begin{aligned} \mathbf{H}_{lm} &= \frac{c_l c_m}{c_d^2 J_d^2} \left( J_m J_l + c_d (J_d J_{lm} - J_m J_{ld} - J_l J_{dm} + \frac{J_l J_m}{J_d} J_{dd}) \right) \Big|_{\mathbf{x}=\mathbf{c}}; \\ \mathbf{H}_{ll} &= \frac{c_l J_l}{c_d J_d} + \frac{c_l^2}{c_d^2 J_d^2} \left( J_l^2 + c_d (J_d J_{ll} - 2 J_l J_{ld} + \frac{J_l^2}{J_d} J_{dd}) \right) \Big|_{\mathbf{x}=\mathbf{c}}, \end{aligned}$$

where  $l, m < d$  and  $l \neq m$ .

*Proof.* The formula for  $b_0$  comes from [PW02, Theorem 3.5]. (Please note the typo therein: the  $z_d H_d$  should be a  $-z_d H_d$ .) We prove the formula for  $\mathbf{H}$  only.

For easy reading, let  $\widehat{\mathbf{x}} = (x_1, \dots, x_{d-1})$ . Since  $J$  is holomorphic and  $J_d(\mathbf{c}) \neq 0$ , there exists a holomorphic function  $g$  guaranteed by the Implicit Function Theorem such that in some open ball around  $\mathbf{c}$ ,  $J(\widehat{\mathbf{x}}, g(\widehat{\mathbf{x}})) = 0$ . As shown in [PW02],  $\mathbf{H}$  is the Hessian (matrix of second partial derivatives) of

$$\widetilde{f}(\widehat{\theta}) := -\log g(c_1 e^{i\theta_1}, \dots, c_{d-1} e^{i\theta_{d-1}}) + i \sum_{j=1}^{d-1} \frac{n_j}{n_d} \theta_j + \log g(\mathbf{c})$$

evaluated at  $\theta = \mathbf{0}$ .

Now, from the Implicit Function Theorem we also get

$$D_l g(\widehat{\mathbf{x}}) = -\frac{J_l(\widehat{\mathbf{x}}, g(\widehat{\mathbf{x}}))}{J_d(\widehat{\mathbf{x}}, g(\widehat{\mathbf{x}}))} \quad (l < d);$$

$$D_m J_l(\widehat{\mathbf{x}}, g(\widehat{\mathbf{x}})) = J_{lm} + J_{ld} D_m g = J_{lm} - J_{ld} \frac{J_m}{J_d} \quad (l \leq d, m < d).$$

Thus, up to an additive constant,  $D_l \widetilde{f} = -i c_l e^{i\theta_l} g^{-1} J_l / J_d$ , and for  $l, m < d$  with  $l \neq m$  we have

$$\begin{aligned} D_{lm} \widetilde{f} \Big|_{\widehat{\theta}=\mathbf{0}} &= c_l c_m \left( -g^{-2} D_m g \frac{J_l}{J_d} + g^{-1} \frac{J_d D_m J_l - J_l D_m J_d}{J_d^2} \right) \Big|_{\mathbf{x}=\mathbf{c}}; \\ D_{ll} \widetilde{f} \Big|_{\widehat{\theta}=\mathbf{0}} &= c_l g^{-1} \frac{J_l}{J_d} + c_l^2 \left( -g^{-2} D_l g \frac{J_l}{J_d} + g^{-1} \frac{J_d D_l J_l - J_l D_l J_d}{J_d^2} \right) \Big|_{\mathbf{x}=\mathbf{c}}. \end{aligned}$$

The result then follows by plugging in the Implicit Function Theorem equations above and simplifying.  $\square$

Several of our examples involve main diagonal asymptotics for symmetric  $J$ . In this case  $h(J, \mathbf{c})$  simplifies greatly.

**Proposition 3.4.** If  $\text{CRIT}(\mathbf{n})$  is finite,  $\mathbf{n} = (n, \dots, n)$ ,  $J(\mathbf{x})$  is symmetric in  $\mathbf{x}$ , and  $J_d(\mathbf{c}) \neq 0$ , where  $\mathbf{c}$  is the contributing point that lies in the positive orthant of  $\mathbb{R}^d$ , then  $\mathbf{c} = (c, \dots, c)$  for some positive real  $c$ , and

$$h(J, \mathbf{c}) = d \left( 1 + \frac{c}{J_d} (J_{dd} - J_{d1}) \Big|_{\mathbf{x}=\mathbf{c}} \right)^{d-1}.$$

*Proof.* By Theorem 3.1,  $\text{CRIT}(\mathbf{n})$  contains exactly one point  $\mathbf{c}$  that lies in the positive orthant of  $\mathbb{R}^d$ . By the symmetry of the critical equations  $J = 0$  and  $x_i J_i = x_d J_d$  ( $i < d$ ) induced by the symmetry of  $\mathbf{n}$  and  $J$ , any permutation of  $\mathbf{c}$ 's coordinates is also a critical point lying in the positive orthant of  $\mathbb{R}^d$ . Since there can be only one such point,  $\mathbf{c} = (c, \dots, c)$  for some positive real  $c$ .

Since  $J$  is symmetric and  $\mathbf{c} = (c, \dots, c)$ , all first partial derivatives of  $J$  are equal at  $\mathbf{c}$ , all mixed second partial derivatives of  $J$  are equal at  $\mathbf{c}$ , and all diagonal second partial derivatives of  $J$  are equal at  $\mathbf{c}$ . Thus the entries of the matrix of Theorem 3.3 simplify to

$$\begin{aligned} D_{lm} \tilde{f} \Big|_{\hat{\theta}=\mathbf{0}} &= 1 + \frac{c}{J_d} (J_{dd} - J_{d1}) \Big|_{\mathbf{x}=\mathbf{c}} \\ D_{ll} \tilde{f} \Big|_{\hat{\theta}=\mathbf{0}} &= 2 D_{lm} \tilde{f} \Big|_{\hat{\theta}=\mathbf{0}}, \end{aligned}$$

where  $l, m < d$  and  $l \neq m$ , and the determinant simplifies to  $h(J, \mathbf{c}) = a^{d-1}d$ , where  $a = 1 + \frac{c}{J_d} (J_{dd} - J_{d1}) \Big|_{\mathbf{x}=\mathbf{c}}$ .  $\square$

Our new diagonal method is simply the application of Theorems 3.1, 3.2, and 3.3 directly to  $F$ . We illustrate this now with several examples.

**Example 3.5 (Zigzag-free Binary Words).** Consider again the bivariate generating function

$$F(x, y) = \sum_{m,n} f_{mn} x^m y^n = \frac{1 + xy + x^2 y^2}{1 - x - y + xy - x^2 y^2}$$

of Section 2. To compute the asymptotics of  $f_{nn}$  using the new diagonal method, let  $\mathbf{n} = (n, n)$  and  $J = 1 - x - y + xy - x^2 y^2$ . Then  $\text{CRIT}(\mathbf{n})$ , the solution set of  $J = 0$  and  $nxJ_x = nyJ_y$ , comprises  $(1/\phi, 1/\phi)$ ,  $(-\phi, -\phi)$ ,  $(1 \pm i\sqrt{3})/2$ , and  $(1 \pm i\sqrt{3})/2$ , where  $\phi = (1 + \sqrt{5})/2$ . By Theorem 3.1,  $\mathbf{c} := (1/\phi, 1/\phi)$  is a contributing point and the only such point, since none of the other critical points lie on the same torus as  $\mathbf{c}$ . It is also a smooth point. By Proposition 3.4,  $h(J, \mathbf{c}) = 4/(-5 + 3\sqrt{5})$ , and by Theorems 3.2 and 3.3,

$$f_{nn} \sim \phi^{2n} \frac{2}{\sqrt{\sqrt{5}\pi n}}.$$

This agrees with our answer in Section 2 since  $\phi^2 = 2/(3 - \sqrt{5})$ .  $\triangleleft$

**Example 3.6 (Lattice Paths).** Consider again the bivariate generating function

$$F(x, y) = \frac{1}{1 - x - y - xy},$$

of Section 2. To compute the asymptotics of  $f_{an, bn}$  using the new diagonal method, let  $\mathbf{n} = (an, bn)$  and  $J = 1 - x - y - xy$ . Then  $\text{CRIT}(\mathbf{n})$ , the solution set of  $J = 0$  and  $bnxJ_x = anyJ_y$ , comprises  $(\frac{-a+L}{b}, \frac{-b+L}{a})$ , where  $L = \sqrt{a^2 + b^2}$ . By Theorem 3.1,  $\mathbf{c} := (\frac{-a+L}{b}, \frac{-b+L}{a})$  is the only contributing point. It is also a smooth point. By Theorem 3.3,  $h(J, \mathbf{c}) = \frac{-2(b-L)a(a^2+b^2-aL)}{(a-L)^2(a-b+L)^2}$ , and by Theorem 3.2,

$$f_{an, bn} \sim \left( \frac{-a+L}{b} \right)^{-an} \left( \frac{-b+L}{a} \right)^{-bn} \sqrt{\frac{ab}{L(a+b-L)^2 2\pi n}}.$$

Here, where the standard diagonal method failed, the new diagonal method allowed us to compute the asymptotics quite easily (with the help of Maple). This success comes from a general attribute of the new diagonal method: the parameters  $a$  and  $b$  appear as factors instead of as powers in the equations we need to solve.  $\triangleleft$

**Example 3.7 (*Ternary Words*).** Consider again the trivariate generating function

$$F(x, y, z) = \frac{1}{1 - x - y - z}$$

of Section 2. To compute the asymptotics of  $f_{an, bn, cn}$ , let  $\mathbf{n} = (an, bn, cn)$  and  $J = 1 - x - y - z$ . Then  $\text{CRIT}(\mathbf{n})$ , the solution set of  $J = 0$ ,  $cnxJ_x = anzJ_z$ , and  $cnyJ_y = bnzJ_z$  comprises  $\mathbf{c} := (a + b + c)^{-1}(a, b, c)$ . By Theorem 3.1,  $\mathbf{c}$  is the only contributing point. It is also a smooth point. By Proposition 3.4,  $h(J, \mathbf{c}) = ab(a + b + c)/c^3$ , and by Theorems 3.2 and 3.3,

$$f_{an, bn, cn} \sim \frac{(a + b + c)^{(a+b+c)n}}{a^{an}b^{bn}c^{cn}} \sqrt{\frac{a + b + c}{abc}} \frac{1}{2\pi n},$$

in agreement with Stirling's formula.

Here, where the standard diagonal method failed because of the nonalgebraic character of the diagonal generating function, the new diagonal method gave the answer easily, even off the main diagonal.  $\triangleleft$

Rather than multiply examples unnecessarily, we finish with an example involving an arbitrary number of variables. Many more examples of multivariate asymptotics using similar methods can be found in [PW].

**Example 3.8 (*Alignments*).** In a problem from computational biology, one is given a finite set of strings of various lengths over some finite alphabet and needs to count the number of ways of inserting blanks into these strings to make them all of equal length. The strings represent genetic sequences, and inserting blanks represents aligning genetically similar segments among the sequences.

More specifically and abstractly, a  $(d, r_1, \dots, r_d)$ -*alignment* is a table of  $d$   $\{0, 1\}$ -strings, one in each row, such that the sum of the entries of row  $j$  equals  $r_j$  ( $1 \leq j \leq d$ ) and no column contains all zeros. (The ones correspond to letters of the genetic sequence and the zeros to blanks.) As shown in [GHOW90], the generating function for the number of  $(d, \cdot)$ -alignments is

$$F(x_1, \dots, x_d) = \frac{1}{2 - \prod_{i=1}^d (1 + x_i)}.$$

To compute the asymptotics of  $f_{n, \dots, n}$  (the case where all rows have the same sum; the genetic sequences all have the same length) using the new diagonal method, let  $\mathbf{n} = (n, \dots, n)$ ,  $I = 1/2$  and  $J = 1 - (1/2) \prod_{i=1}^d (1 + x_i)$ . Then  $\text{CRIT}(\mathbf{n})$ , the solution set of  $J = 0$  and  $nx_iJ_i = nx_dJ_d$  ( $i < d$ ) comprises  $(2^{1/d}\omega_i - 1, \dots, 2^{1/d}\omega_i - 1)$ , where the  $\omega_i$  ( $i < d$ ) are the  $d$ th roots of unity. By Theorem 3.1,  $\mathbf{c} := (2^{1/d} - 1, \dots, 2^{1/d} - 1)$  is the only contributing point. It is also a smooth point. By Proposition 3.4,  $h(J, \mathbf{c}) = d2^{(1-d)/d}$ , and

by Theorems 3.2 and 3.3,

$$f_{n\dots n} \sim (2^{1/d} - 1)^{-dn} \frac{1}{(2^{1/d} - 1)2^{(d^2-1)/2d} \sqrt{d(\pi n)^{d-1}}},$$

in agreement with [GHOW90]. (Please note the typo therein: the  $2^{(k^2-1)/2k}$  in the formula on page 139 should be a  $2^{(1-k^2)/2k}$ ).

A more important case for biological applications is to count alignments whose minimum block size is bounded below by a constant. A *block* of size  $b$  in a  $(d, \cdot)$ -alignment is a  $d \times b$  submatrix of the alignment with contiguous columns, none of which contain a zero. As shown in [RT], the generating function for  $(d, \cdot)$ -alignments with blocks of size at least  $b$  is

$$\begin{aligned} G(x_1, \dots, x_d) &= \frac{1 - t + t^b}{1 + (1 - p)(1 - t + t^b) - t^2 + t^{b+1}} \\ &= \frac{1 - t + t^b}{1 - t - (p - 1 - t)(1 - t + t^b)} \end{aligned}$$

where  $p = \prod_{i=1}^d (1 + x_i)$  and  $t = \prod_{i=1}^d x_i$ . Note that when  $b = 1$ ,  $G = F$ , as expected.

To compute the asymptotics of  $g_{n,\dots,n}$  (which to our knowledge has only been done up till now for  $d = 2$  via the standard diagonal method [RT]) let  $\mathbf{n} = (n, \dots, n)$  and  $J = 1 - t - (p - 1 - t)(1 - t + t^b)$ . Then  $\text{CRIT}(\mathbf{n})$  comprises all vectors  $x\mathbf{1}$  such that  $j(x) := J(x\mathbf{1}) = 0$  (as seen from the equations  $nx_i J_i = nx_d J_d$ ). Let  $\mathbf{c} := c\mathbf{1}$  be the critical point in the positive orthant of  $\mathbb{R}^d$  (so  $c$  is the unique positive solution of  $j(x) = 0$ ) and note that  $c < 1$ , because  $j(x) < 0$  for  $x \geq 1$ . By Theorem 3.1,  $\mathbf{c} \in \text{CONTRIB}(\mathbf{n})$  and all other points of  $\text{CONTRIB}(\mathbf{n})$  lie on the same torus as  $\mathbf{c}$ . Since

$$G(x_1, \dots, x_d) = \frac{1 + \frac{t^b}{1-t}}{1 - (p - 1 - t)(1 + \frac{t^b}{1-t})},$$

on the open polydisk  $\{\mathbf{x} \in \mathbb{C}^d : \forall i |x_i| < 1\}$ , since this polydisk contains the torus of  $\mathbf{c}$ , and since  $(p - 1 - t)(1 + \frac{t^b}{1-t})$  is an aperiodic power series with nonnegative coefficients, we can apply Theorem 3.1 to conclude that  $\mathbf{c}$  is the only contributing point. It is also a smooth point, as is seen most easily from the form  $J = 1 - (p - 1 - t)(1 + \frac{t^b}{1-t})$ . Thus by Theorem 3.2,

$$g_{n\dots n} \sim c^{-dn} b_0 n^{(1-d)/2},$$

where  $b_0$  is the constant given in Theorem 3.3. Note that both  $c$  and  $b_0$  depend on  $b$ .  $\triangleleft$

## References

- [BK86] Egbert Brieskorn and Horst Knörrer, *Plane algebraic curves*, Birkhäuser Verlag, Basel, 1986, Translated from the German by John Stillwell. MR886476 (88a:14001)

- [BP] Yuliy Baryshnikov and Robin Pemantle, *Convolutions of inverse linear functions via multivariate residues*, preprint available at <http://www.math.upenn.edu/~pemantle/papers/Preprints/hyperplanes.pdf>.
- [FS] Phillipe Flajolet and Robert Sedgewick, *Analytic combinatorics*, in preparation, preprint available at <http://algo.inria.fr/flajolet/Publications/book061023.pdf>.
- [Fur67] Harry Furstenberg, *Algebraic functions over finite fields*, J. Algebra **7** (1967), 271–277. MR MR0215820 (35 #6655)
- [GHOW90] J. R. Griggs, P. Hanlon, A. M. Odlyzko, and M. S. Waterman, *On the number of alignments of  $k$  sequences*, Graphs Combin. **6** (1990), no. 2, 133–146. MR MR1073683 (91m:92022)
- [HK71] M. L. J. Hautus and D. A. Klarner, *The diagonal of a double power series*, Duke Math. J. **38** (1971), 229–235. MR MR0276641 (43 #2383)
- [Lip88] L. Lipshitz, *The diagonal of a  $D$ -finite power series is  $D$ -finite*, J. Algebra **113** (1988), no. 2, 373–378. MR MR929767 (89c:13027)
- [MZS04] Emanuele Munarini and Norma Zagaglia Salvi, *Binary strings without zigzags*, Sémin. Lothar. Combin. **49** (2002/04), Art. B49h, 15 pp. (electronic). MR MR2050271 (2005a:05011)
- [PW] Robin Pemantle and Mark C. Wilson, *Twenty combinatorial examples of asymptotics derived from multivariate generating functions*, submitted, preprint available at <http://arxiv.org/pdf/math.CO/0512548>.
- [PW02] ———, *Asymptotics of multivariate sequences. I. Smooth points of the singular variety*, J. Combin. Theory Ser. A **97** (2002), no. 1, 129–161. MR MR1879131 (2003a:05015)
- [PW04] ———, *Asymptotics of multivariate sequences. II. Multiple points of the singular variety*, Combin. Probab. Comput. **13** (2004), no. 4-5, 735–761. MR MR2095981 (2005i:05008)
- [RT] M. Régnier and F. Tahi, *Generating functions in computational biology*, preprint available at <http://algo.inria.fr/regnier/publis/ReTa04.ps>.
- [Sta99] Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin. MR MR1676282 (2000k:05026)