











A new approach to asymptotics of Maclaurin coefficients of algebraic functions

Alexander Raichev Mark C. Wilson

Department of Computer Science University of Auckland Private Bag 92019 Auckland New Zealand

CDMTCS-322 April 2008

Centre for Discrete Mathematics and Theoretical Computer Science

A NEW APPROACH TO ASYMPTOTICS OF MACLAURIN COEFFICIENTS OF ALGEBRAIC FUNCTIONS

ALEXANDER RAICHEV AND MARK C. WILSON

ABSTRACT. We propose a general method for deriving asymptotics of the Maclaurin series coefficients of algebraic functions that is based on a procedure of K. V. Safonov and multivariate singularity analysis. We test the feasibility of this this approach by experimenting on several examples.

1. INTRODUCTION

Let \mathbb{N}_+ denote the set of (strictly) positive natural numbers. For $m \in \mathbb{N}_+$, $\beta \in \mathbb{N}^m$, and $x \in \mathbb{C}^m$, let x_i denote component *i* of *x*, let $\beta! = \beta_1! \cdots \beta_m!$, $n\beta = (n\beta_1, \ldots, n\beta_m)$, $x^\beta = x_1^{\beta_1} \cdots x_m^{\beta_m}$, and $\partial^\beta = \partial_1^{\beta_1} \cdots \partial_m^{\beta_m}$, where ∂_j is partial differentiation with respect to coordinate *j*. For a domain $\Xi \subseteq \mathbb{C}^m$ (an open connected set) containing the origin and a holomorphic function $f: \Xi \to \mathbb{C}$, let $f_\beta = \partial^\beta f(0)/\beta!$ for $\beta \in \mathbb{N}^m$, the coefficients of the Maclaurin series for *f*.

In [Saf00] K. V. Safonov proved the following theorem.

Theorem 1.1. If $f :\subset \mathbb{C}^{d-1} \to \mathbb{C}$ is an algebraic function holomorphic in a neighborhood of the origin and every term of the Maclaurin series of f(x) contains a factor of x_1 , then there exists a rational function $F :\subset \mathbb{C}^d \to \mathbb{C}$ holomorphic in a neighborhood of the origin and a matrix $M \in GL_{d-1}(\mathbb{Z})$ with nonnegative entries such that the following relation between the Maclaurin series coefficients of f and F holds:

$$f_{\beta} = F_{\gamma,\gamma_1},$$

where $\beta \in \mathbb{N}^{d-1}$ and $\gamma = \beta M$.

Loosely speaking, every (d-1)-variate algebraic power series is a generalized diagonal of a *d*-variate rational power series. Moreover, Safonov gave a constructive procedure for finding *F* given an annihilating polynomial for *f*.

In this report we propose a new approach to deriving asymptotics of Maclaurin series coefficients of algebraic functions f, namely, (1) apply Safonov's procedure to f to get a multivariate rational function F, then (2) apply multivariate singularity analysis to F to derive the asymptotics of the Maclaurin series coefficients of F in the direction appropriate to f. In this way we reduce deriving asymptotics of algebraic functions to deriving asymptotics of rational functions. This is promising, because current general methods for the former task exist only in the univariate case (see [FS, Chapter VII] for a summary of these), whereas current general methods for the latter exist in the multivariate case as well (see [PW] for a summary of these).

¹⁹⁹¹ Mathematics Subject Classification. 05A15, 05A16.

Key words and phrases. generating function, multivariate, asymptotics, diagonal, rational, algebraic, resolution of singularities.

We test the feasibility of this approach by experimenting on several specific algebraic functions. In some cases we meet with difficulty because of gaps in the present theory of multivariate singularity analysis for rational functions. We note these gaps and indicate directions for further research.

2. Examples

Let us apply our new approach to several specific algebraic functions, beginning with univariate ones. Their asymptotics are already known, so we can check our answers. We used Maple 11 to do the calculations, and our worksheets are available at http://www.cs.auckland.ac.nz/~raichev/research.html.

A description of Safonov's procedure, which employs resolution of singularities, can be found in [Saf00], and a description of multivariate singularity analysis for rational functions can be found in [ERW] and [PW]. We will assume you are familiar with this material. Warning: [Saf00] contains some confusing passages; see Section 3 for a helpful annotation.

Example 2.1 (Catalan numbers right-shifted). Success.

Consider the algebraic function

$$f(x_1) = \frac{1 - \sqrt{1 - 4x_1}}{2}.$$

It is the generating function of the Catalan numbers multiplied by x_1 , and so its sequence of Maclaurin coefficients is the Catalan sequence prefixed by 0. We seek asymptotics for f_n .

Since the generating function of the Catalan numbers is holomorphic in a neighborhood of the origin, we may apply Safonov's procedure to f. Doing so, we get that $f_n = F_{n,n}$, where

$$F(x_1, x_2) = \frac{x_2(2x_2 - 1)}{x_1 + x_2 - 1}.$$

So the Maclaurin coefficients of f are the diagonal Maclaurin coefficients of F. Notice that this is always the case for univariate f since d = 1 implies M = (1) in Theorem 1.1. Notice also that in this example f is **combinatorial**, that is, its Maclaurin coefficients are all nonnegative, but F is not combinatorial.

To derive asymptotics for f_n , we derive asymptotics for $F_{n\alpha}$ with $\alpha = (1, 1)$. To do this we apply multivariate singularity analysis to F. Letting H equal the denominator of F, we find that $\mathcal{V} = \{x \in \mathbb{C}^2 : H(x) = 0\}$ is globally smooth. There is one critical point of F for α , namely c = (1/2, 1/2), and c is clearly strictly minimal. Thus by the formulas of [ERW] (ERW-formulas),

$$F_{n\alpha} = 4^n \left[\frac{1}{4\sqrt{\pi}n^{3/2}} + \frac{3}{32\sqrt{\pi}n^{5/2}} + O\left(n^{-7/2}\right) \right],$$

as $n \to \infty$, which is correct.

Example 2.2 (*Supertrees*). Failure. Critical points not minimal. Consider the polynomial

 $P(y, x_1) = y^4 - 2y^3 + (1 + 2x_1)y^2 - 2yx_1 + 4x_1^3.$

It is an annihilating polynomial for the combinatorial generating function $f(x_1)$ for supertrees (see [FS] for more details). The first few Maclaurin coefficients of f are known to be 0, 0, 2, 2, 8, 18, 64, 188. Applying Safonov's procedure to f via P we get that $f_n = F_{n,n}$, where

$$F(x_1, x_2) = \frac{2x_1x_2^2(2x_2^5x_1^2 - 3x_2^3x_1 + x_2 + 2x_2^2x_1 - 1)}{x_2^5x_1^2 - 2x_2^3x_1 + x_2 + 2x_2^2x_1 - 2 + 4x_1}.$$

To derive asymptotics for f_n , we derive asymptotics of $F_{n\alpha}$ where $\alpha = (1, 1)$. Letting H equal the denominator of F, we find that \mathcal{V} is globally smooth. There are three critical points of F for α , namely $(3/16 - (1/16)\sqrt{5}, 1 + \sqrt{5}), (3/16 + (1/16)\sqrt{5}, 1 - \sqrt{5}), and <math>(1/8, 2)$. However, none of these points is minimal. To see this, for each of these points q we compute the first nonzero term of the Maclaurin series of \tilde{g} (which depends on q; see [ERW] for a definition of \tilde{h} and \tilde{g}) and get at^4 with a < 0. Thus $\Re \tilde{g} < 0$ near 0, hence $|\tilde{h}(t)| < |q_2|$ near 0, hence there exists a point $q' \in \mathcal{V}$ such that $|q'_1| = |q_1|$ and $|q'_2| < |q'_2|$, hence q is not minimal. So the ERW-formulas are not guaranteed to work.

Plugging c = (1/8, 2) into the formulas anyway we get

$$F_{n\alpha} = 4^n \left[\frac{1 - i}{8\Gamma(3/4)n^{5/4}} + O\left(n^{-7/4}\right) \right]$$

which is nonreal and thus incorrect since $f_n \ge 0$. The correct answer is

$$F_{n\alpha} = 4^n \left[\frac{1}{8\Gamma(3/4)n^{5/4}} + O\left(n^{-7/4}\right) \right]$$

(see [FS, Chapter VI] for instance).

Example 2.3 (*Basic univariate*). Failure. No critical points.

Consider the simple combinatorial algebraic function

$$f(x_1) = \frac{x_1}{\sqrt{1-x_1}}.$$

Applying Safonov's procedure to f we get that $f_n = F_{n,n}$, where

$$F(x_1, x_2) = \frac{x_2 x_1 (-3x_2 + 3x_2^2 x_1 - 2 + 2x_2 x_1 + x_1 - 2x_2^2 + 2x_2^3 x_1)}{-x_2 + x_2^2 x_1 - 2 + 2x_2 x_1 + x_1}$$

To derive asymptotics for f_n we derive asymptotics for $F_{n\alpha}$ where $\alpha = (1, 1)$. Letting H equal the denominator of F, we find that \mathcal{V} is globally smooth. However, there are no critical points of F for α . Thus we can not apply the ERW-formulas.

We try to get around this as follows. When $\alpha = (1, \lambda)$ for $0 < \lambda < 1$, there are two critical points of F, namely, $\left(\frac{1}{8}\lambda^2 + (\frac{1}{8}\lambda - \frac{3}{8})\sqrt{\lambda^2 - 10\lambda + 9} - \lambda + \frac{7}{8}, \frac{3-3\lambda + \sqrt{\lambda^2 - 10\lambda + 9}}{2(\lambda - 1)}\right)$ and $\left(\frac{1}{8}\lambda^2 - (\frac{1}{8}\lambda - \frac{3}{8})\sqrt{\lambda^2 - 10\lambda + 9} - \lambda + \frac{7}{8}, \frac{3-3\lambda - \sqrt{\lambda^2 - 10\lambda + 9}}{2(\lambda - 1)}\right)$. Plugging the second and first critical points into the ERW-formulas and taking the limit as $\lambda \to 1^-$ we get

$$F_{n\alpha} = \frac{1}{\sqrt{2\pi n}} + \infty,$$

and its negative, respectively. So both expansions are clearly incorrect, but the leading term of the first is curiously close to that of the correct expansion, which is

$$F_{n\alpha} = \frac{1}{\sqrt{\pi n}} + O\left(n^{-3/2}\right)$$

(see [FS, Chapter VI] for instance).

Alternatively, searching for critical points of F for $\alpha = (1, 1)$ in \mathbb{CP}^2 yields two points, namely (0, 1, 0) and (0, 0, 1) in homogeneous coordinates. However, mapping these back down to \mathbb{C}^2 eventually leads to a trivariate generating function with cone point singularities, a type of singularity for which there is presently no multivariate singularity analysis worked out.

Example 2.4 (*Basic bivariate*). Success.

Consider the bivariate algebraic function

$$f(x_1, x_2) = x_1 \sqrt{1 - x_1 - x_2},$$

which appears in [Saf00, Example 2]. Following Safonov's procedure we get that $f_{n\beta} = F_{n\beta M}$ where $M = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and

$$F(x_1, x_2, x_3) = \frac{x_3 x_1 (3x_3 + 2 + x_1 + x_2 x_1 + 2x_3^2)}{x_3 + 2 + x_1 + x_2 x_1}$$

Letting H equal the denominator of F, we find that \mathcal{V} is globally smooth. There is one critical point of F for βM , namely $(\frac{-\beta_1}{\beta_1+\beta_2}, \frac{\beta_2}{\beta_1}, -1)$. Since F is not combinatorial, we have no shortcuts for checking whether or not this point is minimal. Proceeding anyway and plugging the critical point into the ERW-formulas we get

$$F_{n\beta M} = -\left[\left(\frac{\beta_1 + \beta_2}{\beta_1}\right)^{\beta_1 + \beta_2} \left(\frac{\beta_1}{\beta_2}\right)^{\beta_2}\right]^n \left[\frac{\beta_1}{\sqrt{8\pi(\beta_1 + \beta_2)^2\sqrt{\beta_1\beta_2}n^2}} + O\left(n^{-3}\right)\right],$$

as $n \to \infty$ uniformly as βM normalized to length 1 varies within a compact set in \mathbb{QP}^2 . This expansion agrees with experimental checks.

Example 2.5 (Another basic bivariate). Failure. No critical points.

Consider the bivariate algebraic function

$$f(x_1, x_2) = \frac{x_1}{\sqrt{1 - x_1 - x_2}}$$

Following Safonov's procedure we get that $f_{n\beta} = F_{n\beta M}$ where $M = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$ and

$$F(x_1, x_2, x_3) = \frac{x_3 x_1 (-3 x_3 + 3 x_3^2 x_1 + 3 x_3^2 x_1 x_2 - 2 + 2 x_3 x_1 + 2 x_3 x_1 x_2 + x_1 + x_1 x_2 - 2 x_3^2 + 2 x_3^3 x_1 + 2 x_3^3 x_1 x_2)}{-x_3 + x_3^2 x_1 + x_3^2 x_1 x_2 - 2 + 2 x_3 x_1 + 2 x_3 x_1 x_2 + x_1 + x_1 x_2}.$$

Letting H equal the denominator of F, we find that \mathcal{V} is globally smooth. However there are no critical points of F for βM . Thus it is unclear how best to proceed in deriving asymptotics. It seems we should first understand the no-critical-points phenomenon in the simpler bivariate setting of Example 2.3.

3. Comments and Questions

As we see, to implement our new approach for deriving asymptotics of Maclaurin coefficients of algebraic functions in full generality, several gaps in the theory of multivariate singularity analysis of rational functions need to be filled, namely,

- How does one check for minimality in the non-combinatorial case?
- How does one handle non-minimal critical points?
- How does one handle the case of no critical points?

4

Thus more research is required.

We end with an observation on Safonov's procedure and a short annotation of [Saf00].

Proposition 3.1. Let F be the rational function from Theorem 1.1. At a critical point of F for $\alpha := (m_1, \ldots, m_{d-1}, m_1) \in \mathbb{N}^d_+$ whose last component is nonzero, the numerator of F vanishes.

Proof. From Safonov's procedure, F(x) has the form "polynomial in $x + (power of x) \cdot R(x)$ ", where

$$R(\hat{x}, y) = \frac{y^2}{k} \frac{(\partial_d P)(x_1 y, x_2, \dots, x_{d-1}, y)}{P(x_1 y, x_2, \dots, x_{d-1}, y)},$$

 $y = x_d$ (for easy reading), $k \in \mathbb{N}_+$, P is a polynomial, and the variable substitution is understood to occur after the partial differentiation. By definition, the denominator of a rational function vanishes at a critical point. Thus by the specific form of F above, it suffices to prove the proposition for the rational function $R(\hat{x}, y)$.

Let J be the denominator of R. Among the critical equations of R for α are

$$J(\hat{x}, y) = 0$$

$$\alpha_1^{-1} x_1 \partial_1 J(\hat{x}, y) = \alpha_d^{-1} y \partial_d J(\hat{x}, y),$$

that is,

 $P(x_1y, x_2, \dots, x_{d-1}, y) = 0$

 $x_1 y(\partial_1 P)(x_1 y, x_2, \dots, x_{d-1}, y) = y [x_1(\partial_1 P)(x_1 y, x_2, \dots, x_{d-1}, y) + (\partial_d P)(x_1 y, x_2, \dots, x_{d-1}, y)].$ Thus at a critical point where $y \neq 0$, $P(x_1 y, x_2, \dots, x_{d-1}, y) = 0$ and $(\partial_d P)(x_1 y, x_2, \dots, x_{d-1}, y) = 0$, as desired.

Corollary 3.2. For a critical point of F whose last component is nonzero, the first term in the ERW-formulas for the asymptotic expansion of F is zero.

Proof. By [ERW, Proposition 4.3], the first term in the ERW-formulas for the asymptotic expansion of $F_{n\alpha}$ contains a factor of the numerator of F evaluated at the critical point. Thus by the previous proposition this term is zero.

In retrospect we should have expected no critical points in Example 2.3, for in the presence of a critical point the ERW-formulas and Corollary 3.2 predict no $n^{-1/2}$ term in the expansion for $F_{n\alpha}$ whereas the correct expansion does have an $n^{-1/2}$ term.

Annotation to [Saf00]

- page 261, line 4 of the main body of the article: z should be z_1
- page 263, lines 2–5: it appears that Safonov is defining his term 'A-diagonal' here
- 264, 6: \widetilde{R} should be R(w, z)
- 266, 1: 'of the degree q' should be 'of some degree q'
- 266, 7: ζ^{m_n} should be $\zeta_n^{m_n}$ and the remark 'for some $m_1, \ldots, m_n \in \mathbb{N}$ ' should be added
- 267, 12: the asterisks should be to the left of the rightmost parenthesis
- 269, 9: $u(\zeta)$ should be $u(v, \zeta)$
- 269, last line: here Safonov states that there exists such an *l* but does not explain how to find it; it seems one need to look at a few terms of the Maclaurin expansions

of both h and a and find the l that works by trial and error; note that l is needed for the definition of $R^{(2)}$ on page 270

- 270, 2: $\zeta_1^{\nu_{n1}}$ should be $\zeta_1^{\nu_{1n}}$
- 270, 11: $h_1(\zeta)$ should be $h^{(1)}(\zeta)$
- 270, 13: the leftmost occurrence of $R^{(1)}(z_0, z)$ should be $R^{(2)}(z_0, z)$
- 270, 13: $z_n^{l_2}$ should be $z_n^{l_n}$
- 271, 7: one of the (pk)!s should be a (qk)!

References

- [ERW] A. F. M. ter Elst, Alexander Raichev, and Mark C. Wilson, Asymptotics of coefficients of multivariate generating functions: improvements for smooth points, Submitted, preprint available at http://arxiv.org/.
- [FS] Phillipe Flajolet and Robert Sedgewick, *Analytic combinatorics*, In preparation, preprint available at http://algo.inria.fr/flajolet/Publications/book061023.pdf.
- [PW] Robin Pemantle and Mark C. Wilson, Twenty combinatorial examples of asymptotics derived from multivariate generating functions, to appear in SIAM Review, preprint available at http: //arxiv.org/pdf/math.CO/0512548.
- [Saf00] K. V. Safonov, On power series of algebraic and rational functions in \mathbb{C}^n , J. Math. Anal. Appl. **243** (2000), no. 2, 261–277. MR MR1741523 (2001f:32001)

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF AUCKLAND, PRIVATE BAG 92019, AUCKLAND, NEW ZEALAND *E-mail address*: raichev@cs.auckland.ac.nz

DEPARTMENT OF COMPUTER SCIENCE, UNIVERSITY OF AUCKLAND, PRIVATE BAG 92019, AUCKLAND, NEW ZEALAND *E-mail address:* mcw@cs.auckland.ac.nz