

Sobel filter example

- Compute G_x and G_y, gradients of the image performing the convolution of Sobel kernels with the image
- Use zero-padding to extend the image

0	0	10	10	10
0	0	10	10	10
0	0	10	10	10
0	0	10	10	10
0	0	10	10	10

y

1	0	-1
2	0	-2
1	0	-1

h_x

-1	-2	-1
0	0	0
1	2	1

h_y

G_x

0	30	30	0	-30
0	40	40	0	-40
0	40	40	0	-40
0	40	40	0	-40
0	30	30	0	-30

G_y

0	-10	-30	-40	-30
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	10	30	40	30

1

Sobel filter example

- Compute G_x and G_y, gradients of the image performing the convolution of Sobel kernels with the image
- Use border values to extend the image

0	0	10	10	10
0	0	10	10	10
0	0	10	10	10
0	0	10	10	10
0	0	10	10	10

y

1	0	-1
2	0	-2
1	0	-1

h_x

-1	-2	-1
0	0	0
1	2	1

h_y

G_x

0	40	40	0	0
0	40	40	0	0
0	40	40	0	0
0	40	40	0	0
0	40	40	0	0

G_y

0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0

$$\Theta = \arctan\left(\frac{G_y}{G_x}\right)$$

0	0		
0	0		
0	0		
0	0		
0	0		

2

Sobel filter example

- Compute G_x and G_y, gradients of the image performing the convolution of Sobel kernels with the image
- Use border values to extend the image

0	0	0	0	0
0	0	0	0	0
10	10	10	10	10
10	10	10	10	10
10	10	10	10	10

y

1	0	-1
2	0	-2
0	0	0
1	0	-1

h_x

-1	-2	-1
0	0	0
1	2	1

h_y

0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0

0	0	0	0	0
-40	-40	-40	-40	-40
-40	-40	-40	-40	-40
0	0	0	0	0
0	0	0	0	0

-π/2	-π/2	-π/2	-π/2	-π/2
-π/2	-π/2	-π/2	-π/2	-π/2

3

$$\Theta = \arctan\left(\frac{G_y}{G_x}\right)$$

Sobel filter example

- Compute G_x and G_y, gradients of the image performing the convolution of Sobel kernels with the image
- Use border values to extend the image

0	0	0	0	10
0	0	0	10	10
0	0	10	10	10
0	10	10	10	10
10	10	10	10	10

y

1	0	-1
2	0	-2
0	0	0
1	0	-1

h_x

-1	-2	-1
0	0	0
1	2	1

h_y

0	0	10	40	30
0	10	30	30	10
10	30	30	10	0
20	30	10	0	0
10	10	0	0	0

0	0	-10	-20	-10
0	-10	-30	-30	-10
-10	-30	-30	-10	0
-40	-30	-10	0	0
-30	-10	0	0	0

		-π/4	-0.15π	-0.10π
		-π/4	-π/4	-π/4
-π/4	-π/4	-π/4	-π/4	
-0.35π	-π/4	-π/4		
-0.40π	-π/4			

4

Laplacian example

- Compute the convolution of image I with the Laplacian kernel
- Use border values to extend the image

0	0	0	0	10
0	0	0	10	10
0	0	10	10	10
0	10	10	10	10
10	10	10	10	10

1	1	1
1	-8	1
1	1	1

Laplacian

0	0	10	40	-20
0	10	30	-30	-10
10	30	-30	-10	0
40	-30	10	0	0
-20	-10	0	0	0

Laplacian*I

5

Erosion example

- Erode the image below using kernels S1 and S2
- Use border values to extend the image

0	0	0	0	1
0	0	0	1	1
0	0	1	1	1
0	1	0	0	1
1	1	1	1	1

1	1	1
1	1	1
1	1	1

S₁

fes1

0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0

0	1	0
1	1	1
0	1	0

S₂

fes2

0	0	0	0	0
0	0	0	0	1
0	0	0	0	1
0	0	0	0	0
0	1	0	0	1

6

Dilation example

- Erode the image below using kernels S1 and S2
- Use border values to extend the image

		$f \oplus S_1$																										
		<table border="1"> <tbody> <tr><td>0</td><td>0</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>0</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> </tbody> </table>	0	0	1	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
0	0	1	1	1																								
0	1	1	1	1																								
1	1	1	1	1																								
1	1	1	1	1																								
1	1	1	1	1																								
y	x																											
		$f \oplus S_2$																										
		<table border="1"> <tbody> <tr><td>0</td><td>0</td><td>0</td><td>1</td><td>1</td></tr> <tr><td>0</td><td>0</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>0</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> </tbody> </table>	0	0	0	1	1	0	0	1	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
0	0	0	1	1																								
0	0	1	1	1																								
0	1	1	1	1																								
1	1	1	1	1																								
1	1	1	1	1																								
		S_1	S_2																									
			7																									

Dilation-Erosion example

- Dilate then erode the image below using kernels S2
- Use border values to extend the image

		$f \oplus S_2$																										
		<table border="1"> <tbody> <tr><td>0</td><td>0</td><td>0</td><td>1</td><td>1</td></tr> <tr><td>0</td><td>0</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>0</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> </tbody> </table>	0	0	0	1	1	0	0	1	1	1	0	1	1	1	1	1	1	1	1	1	1	1	1	1	1	
0	0	0	1	1																								
0	0	1	1	1																								
0	1	1	1	1																								
1	1	1	1	1																								
1	1	1	1	1																								
y	x																											
		$(f \oplus S_2) \ominus S_2$																										
		<table border="1"> <tbody> <tr><td>0</td><td>0</td><td>0</td><td>0</td><td>1</td></tr> <tr><td>0</td><td>0</td><td>0</td><td>1</td><td>1</td></tr> <tr><td>0</td><td>0</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>0</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> <tr><td>1</td><td>1</td><td>1</td><td>1</td><td>1</td></tr> </tbody> </table>	0	0	0	0	1	0	0	0	1	1	0	0	1	1	1	0	1	1	1	1	1	1	1	1	1	
0	0	0	0	1																								
0	0	0	1	1																								
0	0	1	1	1																								
0	1	1	1	1																								
1	1	1	1	1																								
		S_2																										
			8																									