




**CDMTCS
Research
Report
Series**




**Informal Constructive
Reverse Mathematics**





H. Ishihara

Japan Advanced Institute of Science and
Technology, Japan



CDMTCS-229
January 2004



Centre for Discrete Mathematics and
Theoretical Computer Science

Informal Constructive Reverse Mathematics

Hajime Ishihara*

January 22, 2004

Bishop's constructive mathematics (BISH) [1, 2, 7, 27] is an informal mathematics using intuitionistic logic and assuming some function existence axioms:

the axiom of countable choice

$$\forall n \in \mathbb{N} \exists x \in X A(n, x) \rightarrow \exists f \in X^{\mathbb{N}} \forall n \in \mathbb{N} A(n, f(n));$$

the axiom of dependent choice

$$\begin{aligned} \forall x \in X \exists y \in X A(x, y) \rightarrow \\ \forall x \in X \exists f \in X^{\mathbb{N}} (f(0) = x \wedge \forall n \in \mathbb{N} A(f(n), f(n+1))); \end{aligned}$$

the axiom of unique choice

$$\forall x \in X \exists! y \in Y A(x, y) \rightarrow \exists f \in Y^X \forall x \in X A(x, f(x)).$$

It is a core of the varieties of mathematics in the sense that it can be extended not only to intuitionistic mathematics (INT) by adding the principle of continuous choice and the fan theorem [7, 8, 9, 10, 27], and constructive recursive mathematics (RUSS) by adding Markov's principle and the extended Church's thesis [7, 22, 27], but also to classical mathematics (CLASS) practised by most mathematicians today by adding the principle of the excluded middle and the full axiom of choice.

Bishop's constructive (forward) mathematicians have been making every effort, for a given classical theorem A , to find its constructive substitute A' such that

$$\text{BISH} \vdash A' \text{ and } \text{CLASS} \vdash A \leftrightarrow A' \text{ }^1.$$

Of course, it happens that sometimes we can take A' as A ; for examples we can prove the following classical theorems in BISH.

*School of Information Science, Japan Advanced Institute of Science and Technology, Tatsunokuchi, Ishikawa 923-1292, Japan

¹In general, we may find more than one such A' , say A'_1, \dots, A'_n , and in this case, we try to find A'_k such that $\text{BISH} \vdash A'_k \rightarrow A'_i$ for all $i = 1, \dots, n$. In some cases, we have to be contented ourselves with A' such that $\text{BISH} \vdash A'$, $\text{CLASS} \vdash A \rightarrow A'$, and it is strong enough for applications.

Theorem 1 (The completeness of \mathbb{R}). *Every Cauchy sequence of real numbers converges.*

Theorem 2 (The constructive compactness of $[0, 1]$). *$[0, 1]$ is totally bounded and complete.*

Theorem 3 (The Baire category theorem). *The intersection of a sequence of dense open subsets of a complete metric space is dense.*

When A and A' are not equivalent in BISH, we also try to show that A does not admit a constructive proof by giving a **Brouwerian counterexample** to A such that

$$\text{BISH} \vdash A \rightarrow P \text{ and } \text{BISH} \not\vdash P^2$$

for some principle P . The constructive compactness of $[0, 1]$ is classically equivalent to the following special case of **the Bolzano-Weierstraß theorem**:

The sequential compactness of $[0, 1]$. *Every sequence of $[0, 1]$ has a convergent subsequence.*

But it is well known that the sequential compactness of $[0, 1]$ entails in BISH **the limited principle of omniscience (LPO)**:

$$\forall \alpha \in \mathbb{N}^{\mathbb{N}} [\exists n (\alpha(n) \neq 0) \vee \neg \exists n (\alpha(n) \neq 0)],$$

which is an instance of **the Principle of the Excluded Middle** $P \vee \neg P$, and false both in INT and in RUSS [7, 27].

Mandelkern [23] showed its converse and proved the equivalence between the Bolzano-Weierstraß theorem and LPO in BISH, which led the subsequent research of informal constructive reverse mathematics aiming at finding a logical principle P such that

$$\text{BISH} \vdash A \leftrightarrow P,$$

not only for a theorem A in CLASS but also for a theorem A in INT and in RUSS even if it is inconsistent with CLASS. This is possible because CLASS, INT and RUSS are extensions of BISH. In the rest of the paper, we will overview the results in informal constructive reverse mathematics to date; see also Mandelkern [23], Ishihara [11], Bridges, Ishihara and Schuster [4, 5], and Ishihara and Schuster [18] for informal constructive reverse mathematics with various compactness principles; Ishihara [12, 13], Bridges, Ishihara, Schuster and Vîță [6], and Bridges, Ishihara and Schuster [5] with various continuity principles; Ishihara and Schuster [19] with Baire's theorem and its contraposition; see [26] for formal classical reverse mathematics.

The first class of theorems consists of theorems which are equivalent to LPO.

²Since BISH is an informal mathematics, this does not mean formal unprovability, but unacceptability, or at least high dubitation in BISH

Theorem 4. *The following are equivalent in BISH.*

1. LPO.
2. $\forall x \in \mathbb{R}(0 < x \vee \neg(0 < x))$.
3. **The monotone convergence theorem** [23]. *Every bounded monotone sequence of real numbers converges.*
4. **The Bolzano-Weierstraß theorem** [23]. *Every bounded sequence of real numbers has a convergent subsequence.*
5. **The sequential compactness theorem** [18]. *Every compact metric space is sequentially compact.*³
6. **The pseudo Heine-Borel theorem** [25]. *Every sequence of closed sets of a compact metric space with the finite intersection property has nonempty intersection.*

The second class of theorems consists of theorems which are equivalent to **the weak limited principle of omniscience** (WLPO):

$$\forall \alpha \in \mathbb{N}^{\mathbb{N}}[\neg \exists n(\alpha(n) \neq 0) \vee \neg \neg \exists n(\alpha(n) \neq 0)].$$

WLPO is an instance of the Principle of the Excluded Middle, weaker than LPO, and false both in INT and in RUSS [7, 27].

Theorem 5. *The following are equivalent in BISH.*

1. WLPO.
2. $\forall x \in \mathbb{R}(\neg(0 < x) \vee \neg \neg(0 < x))$.
3. **The existence of a discontinuous function** [28]. *A discontinuous function from $\mathbb{N}^{\mathbb{N}}$ into \mathbb{N} exists.*⁴

Theorem 6. *The following are equivalent in BISH.*

1. \neg WLPO.
2. **The nondiscontinuity theorem** [13]. *Every mapping of a complete metric space into a metric space is nondiscontinuous.*⁵

³A metric space is *compact* if it is totally bounded and complete, and *sequentially compact* if every sequence of its elements has a convergent subsequence.

⁴A function f between metric spaces is *discontinuous* if there exist $\delta > 0$ and a sequence $\{x_n\}$ converging to a limit x such that $d(f(x_n), f(x)) \geq \delta$ for all n .

⁵A function f between metric spaces is *nondiscontinuous* if $x_n \rightarrow x$ and $d(f(x_n), f(x)) \geq \delta$ for all n imply $\delta \leq 0$.

The third class of theorems consists of theorems which are equivalent to **the lesser limited principle of omniscience** (LLPO):

$$\forall \alpha \beta \in \mathbb{N}^{\mathbb{N}} [\neg(\exists n(\alpha(n) \neq 0) \wedge \exists n(\beta(n) \neq 0)) \rightarrow \neg \exists n(\alpha(n) \neq 0) \vee \neg \exists n(\beta(n) \neq 0)].$$

LLPO is an instance of **De Morgan's law** $\neg(P \wedge Q) \rightarrow \neg P \vee \neg Q$, weaker than WLPO, and false both in INT and in RUSS [7, 27].

Theorem 7. *The following are equivalent in BISH.*

1. LLPO.
2. $\forall x \in \mathbb{R}(\neg(0 < x) \vee \neg(x < 0))$.
3. $\forall xy \in \mathbb{R}(xy = 0 \rightarrow x = 0 \vee y = 0)$.
4. For all $x, y \in \mathbb{R}$ with $\neg(x < y)$, $\{x, y\}$ is closed subset of \mathbb{R} [14].
5. **The weak König lemma** (WKL) [11]. *Every infinite tree has an infinite path.* ⁶
6. **The intermediate value theorem** [7]. *If $f : [a, b] \rightarrow \mathbb{R}$ is a uniformly continuous function, and y is a real number such that $f(a) < y < f(b)$, then there exists x in $[a, b]$ such that $f(x) = y$.*
7. **The minimum principle** [11, 18]. *Every uniformly continuous real function f on a compact metric space X attains its minimum.* ⁷
8. **The pseudo Heine-Borel theorem for zero sets** ⁸ [11, 18]. *Every sequence of zero sets of a compact metric space with the finite intersection property has nonempty intersection.*
9. **The Hahn-Banach theorem** [11]. *Every bounded linear functional f on a subspace of a separable normed space E , whose kernel is located in E ⁹, has an extension g with $\|f\| = \|g\|$.*

The fourth class of theorems consists of theorems which are equivalent to **the fan theorem** (FAN):

Every detachable bar is uniform. ¹⁰

⁶A (binary) *tree* is a subset T of $\mathbf{2}^{<\mathbb{N}}$ such that it is *detachable* from $\mathbf{2}^{<\mathbb{N}}$ in the sense that for each $n \in \mathbf{2}^{<\mathbb{N}}$, either $n \in T$ or $n \notin T$, and it is closed under restriction, that is if $n \in T$ and $m \preceq n$, then $m \in T$. A tree T is *infinite* if for each k there exists $n \in \mathbf{2}^k$ such that $n \in T$, and $\alpha \in \mathbf{2}^{\mathbb{N}}$ is an *infinite path* in T if $\forall k(\bar{\alpha}k \in T)$.

⁷That is there exists x in X such that $f(x) \leq f(y)$ for all $y \in X$.

⁸A subset S of a metric space X is a *zero set* if there is a pointwise continuous function $f : X \rightarrow \mathbb{R}$ such that $S = \{x \in X \mid f(x) = 0\}$.

⁹A subset S of a metric space X is said to be *located* in X if $d(x, S) := \inf\{d(x, y) : y \in S\}$ exists for each x in X .

¹⁰A subset B of $\mathbf{2}^{<\mathbb{N}}$ is called a *bar* if for each $\alpha \in \mathbf{2}^{\mathbb{N}}$ there exists $n \in \mathbb{N}$ such that $\bar{\alpha}n \in B$. A bar B is *uniform* if there exists k such that for each $\alpha \in \mathbf{2}^{\mathbb{N}}$ $\exists i \leq k(\bar{\alpha}i \in B)$.

FAN is a contrapositive form of WKL, weaker than LLPO [11] and hence than WKL [17], accepted in INT, and false in RUSS [7, 27].

Theorem 8. *The following are equivalent in BISH.*

1. FAN.
2. **The uniform continuity theorem** [27, 29]. *Every pointwise continuous mapping from $2^{\mathbb{N}}$ into \mathbb{N} is uniformly continuous.*
3. **The Heine-Borel theorem for cozero sets**¹¹ [18]. *Every cover of a compact metric space by a sequence of cozero sets has a finite subcover.*

The fifth class of theorems consists of theorems which are equivalent to **Markov's principle** (MP):

$$\forall \alpha \in \mathbb{N}^{\mathbb{N}} [\neg \neg \exists n (\alpha(n) \neq 0) \rightarrow \exists n (\alpha(n) \neq 0)].$$

MP is an instance of **the double negation elimination** $\neg \neg P \rightarrow P$, weaker than LPO, rejected in INT, and accepted in RUSS [7, 27].

Theorem 9. *The following are equivalent in BISH.*

1. MP.
2. $\forall x \in \mathbb{R} (\neg \neg (0 < x) \rightarrow 0 < x)$.
3. **The strong extensionality theorem** [3, 14]. *Every mapping between metric spaces is strongly extensional.*¹²

The sixth class of theorems consists of theorems which are equivalent to **weak Markov's principle** (WMP):

$$\forall \alpha \in \mathbb{N}^{\mathbb{N}} [\forall \beta \in \mathbb{N}^{\mathbb{N}} (\neg \neg \exists n (\beta(n) \neq 0) \vee \neg \neg \exists n (\alpha(n) \neq 0 \wedge \beta(n) = 0) \rightarrow \exists n (\alpha(n) \neq 0)].$$

WMP is weaker than MP, provable both in INT and in RUSS [15, 21].

Theorem 10. *The following are equivalent in BISH.*

1. WMP.
2. $\forall x \in \mathbb{R} [\forall y \in \mathbb{R} (\neg \neg (0 < y) \vee \neg \neg (y < x)) \rightarrow 0 < x]$.
3. **The strong extensionality theorem for complete spaces** [13]. *Every mapping from a complete metric space into a metric space is strongly extensional.*

¹¹A subset S of a metric space X is a *cozero set* if there is a pointwise continuous function $f : X \rightarrow \mathbb{R}$ such that $S = \{x \in X \mid f(x) \neq 0\}$.

¹²A mapping f between metric spaces is *strongly extensional* if $f(x) \neq f(y)$ implies $x \neq y$.

4. **The sequential continuity theorem** [13]. *Every nondiscontinuous mapping from a complete metric space to a metric space is sequentially continuous.*¹³

The seventh class of theorems consists of theorems which are equivalent to **the disjunctive version of Markov's principle** (MP^\vee):

$$\forall \alpha \beta \in \mathbb{N}^{\mathbb{N}} [\neg(\neg \exists n(\alpha(n) \neq 0) \vee \neg \exists n(\beta(n) \neq 0)) \rightarrow \neg \neg \exists n(\alpha(n) \neq 0) \vee \neg \neg \exists n(\beta(n) \neq 0)].$$

MP^\vee is an instance of De Morgan's law, weaker than MP and than LLPO, rejected in INT, accepted in RUSS, and, together with WMP, implies MP [15].

Theorem 11. *The following are equivalent in BISH.*

1. MP^\vee .
2. $\forall x \in \mathbb{R} [\neg \neg(x \neq 0) \rightarrow \neg \neg(0 < x)] \vee \neg \neg(x < 0)$.
3. For all $x, y \in \mathbb{R}$ with $\neg \neg(x < y)$, $\{x, y\}$ is closed subset of \mathbb{R} .

The eighth class of theorems consists of theorems which are equivalent to **the boundedness principle** (BD-N):

Every countable pseudobounded subset of \mathbb{N} is bounded,

where a subset A of \mathbb{N} is said to be *pseudobounded* if for each sequence $\{a_n\}$ in A , $a_n < n$ for all sufficiently large n . BD-N is weaker than LPO and provable both in INT and in RUSS [13].

Theorem 12. *The following are equivalent in BISH.*

1. BD-N.
2. **The pointwise continuity theorem** [13]. *Every sequentially continuous mapping from a separable metric space into a metric space is pointwise continuous.*
3. **Banach's inverse mapping theorem** [16]. *If T is a bounded one-one linear mapping of a separable Banach space E onto a Banach space F , then T^{-1} is bounded.*

References

- [1] E. Bishop, *Foundations of Constructive Mathematics*, McGraw-Hill, New York, 1967.
- [2] E. Bishop and D. Bridges, *Constructive Analysis*, Springer, Berlin, 1985.
- [3] D. Bridges and H. Ishihara, *Linear mappings are fairly well-behaved*, Arch. Math. (Basel) **54** (1990), 558-562.

¹³A mapping f between metric spaces is *sequentially continuous* if $x_n \rightarrow x$ implies $f(x_n) \rightarrow f(x)$.

- [4] D. Bridges, H. Ishihara and P. Schuster, *Sequential compactness in constructive analysis*, Österr. Akad. Wiss. Math.-Natur. Kl. Sitzungsber. II **208** (1999), 159-163.
- [5] D. Bridges, H. Ishihara and P. Schuster, *Compactness and continuity, constructively revisited*, in: J. Bradfield ed., *Computer Science Logic*, Springer, Berlin, Lect. Notes Comput. Sci. **2471** (2002), 89-102.
- [6] D. Bridges, H. Ishihara, P. Schuster and L. Viță, *Apartness continuity implies uniformly sequential continuity*, preprint, 2001.
- [7] D. Bridges and F. Richman, *Varieties of Constructive Mathematics*, Cambridge University Press, Cambridge, 1987.
- [8] L.E.J. Brouwer, *Brouwer's Cambridge Lectures on Intuitionism*, ed. D. van Dalen, Cambridge University Press, Cambridge, 1981.
- [9] M. Dummett, *Elements of Intuitionism*, 2nd ed., Clarendon Press, Oxford, 2000.
- [10] A. Heyting, *Intuitionism, An Introduction*, 3rd rev. ed., North-Holland, Amsterdam, 1971.
- [11] H. Ishihara, *An omniscience principle, the König lemma and the Hahn-Banach theorem*, Z. Math. Logik Grundlagen Math. **36** (1990), 237-240.
- [12] H. Ishihara, *Continuity and nondiscontinuity in constructive mathematics*, J. Symbolic Logic **56** (1991), 1349-1354.
- [13] H. Ishihara, *Continuity properties in constructive mathematics*, J. Symbolic Logic **57** (1992), 557-565.
- [14] H. Ishihara, *An introduction to constructive analysis*, preprint, 1992. (Japanese)
- [15] H. Ishihara, *Markov's principle, Church's thesis and Lindelöf's theorem*, Indag. Math. (N.S.) **4** (1993), 321-325.
- [16] H. Ishihara, *Sequential continuity in constructive mathematics*, In: C.S. Calude, M.J. Dinneen and S. Sburlan eds., *Combinatorics, Computability and Logic*, Proceedings of the Third International Conference on Combinatorics, Computability and Logic, (DMTCS'01) in Constanța Romania, July 2-6, 2001, Springer-Verlag, London, 2001, 5-12.
- [17] H. Ishihara, *WKL implies FAN: a direct proof*, preprint, December 2002.
- [18] H. Ishihara and P. Schuster, *Constructive compactness continued*, preprint, 2002.
- [19] H. Ishihara and P. Schuster, *A note on the schema of weak continuity for numbers*, preprint, 2002.
- [20] K-I Ko, *Complexity Theory of Real Functions*, Birkhäuser, Boston, 1991.
- [21] U. Kohlenbach, *On weak Markov's principle*, MLQ Math. Log. Q. **48** (2002), 59-65.

- [22] B. Kushner, *Lectures on Constructive Mathematical Analysis*, Amer. Math. Soc., Providence, 1984.
- [23] M. Mandelkern, *Limited omniscience and the Bolzano-Weierstrass principle*, Bull. London Math. Soc. **20** (1988), 319-320.
- [24] M.B. Pour-El and J.I. Richards, *Computability in Analysis and Physics*, Springer, Berlin, 1989.
- [25] F. Richman, *personal communication*, May 1999.
- [26] S.G. Simpson, *Subsystems of Second Order Arithmetic*, Springer, Berlin, 1999.
- [27] A. S. Troelstra and D. van Dalen, *Constructivism in Mathematics*, two volumes, North-Holland, Amsterdam, 1988.
- [28] M. van Atten and D. van Dalen, *Arguments for the continuity principle*, Bull. Symbolic Logic **8** (2002), 329-347.
- [29] F. Waaldijk, *On the foundations of constructive mathematics*, preprint, April 2001.
- [30] K. Weihrauch, *Computable Analysis*, Springer, Berlin, 2000.