






**CDMTCS
Research
Report
Series**




**A simple construction of
absolutely disjunctive
Liouville numbers**




Cristian S. Calude
University of Auckland



Ludwig Staiger
Martin-Luther-Universität
Halle-Wittenberg



CDMTCS-505
March 2017



Centre for Discrete Mathematics and
Theoretical Computer Science

A simple construction of absolutely disjunctive Liouville numbers

*Cristian S. Calude**

Department of Computer Science, University of Auckland
Auckland, New Zealand

and

Ludwig Staiger†

Martin-Luther-Universität Halle-Wittenberg
Institut für Informatik
von-Seckendorff-Platz 1, D–06099 Halle (Saale), Germany

Abstract

A disjunctive sequence is an infinite sequence in which every finite string appears as a substring. An absolutely disjunctive number (or lexicon) is a real whose expansion with respect to every base is disjunctive.

In this note we give a simple construction of absolutely disjunctive Liouville numbers (reals which can be “quite closely” approximated by sequences of rationals).

Keywords: Disjunctive sequences, Liouville numbers, computability

1 Introduction

Disjunctivity is a qualitative form of (Borel) normality: normal sequences are disjunctive, but the converse is false. Like normality [[Cs59](#),

*email: cristian@cs.auckland.ac.nz

†email: ludwig.staiger@informatik.uni-halle.de

[Sc60], disjunctivity is not base-invariant (for more details see [He96]).

Jürgensen and Thierrin [JT88] gave a construction of Liouville numbers disjunctive in base b . Highly incomputable Liouville numbers disjunctive to every base have been presented in [St02a, Theorem 15].

The recent construction of a computable absolutely normal Liouville number in [BHS15] yields also computable, absolutely disjunctive Liouville numbers. This construction, however, is based on rather complicated measure-theoretic arguments from [Bu02]. The aim of this note is to present a simple algorithm producing weaker examples, that is, computable Liouville numbers disjunctive to every base.

1.1 Notation

In this section we introduce the notation used throughout the paper. By $\mathbb{N} = \{0, 1, 2, \dots\}$ we denote the set of natural numbers. Its elements will be usually denoted by letters i, \dots, n . The set $A_b = \{0, 1, \dots, b-1\}$, where $b \geq 2$ is a positive integer, is called the b -base; the elements of A_b are called b -digits. By A_b^* we denote the set of all finite strings (words) with ε denoting the empty string; A_b^ω is the set of all infinite sequences (ω -words) over A_b ; ω -words are usually denoted by x, y . The length of a finite or infinite string η over A_b is denoted by $|\eta|$.

For $w \in A_b^*$ and $\eta \in A_b^* \cup A_b^\omega$, $w \cdot \eta$ is their *concatenation*. This concatenation product extends in an obvious way to subsets $L \subseteq A_b^*$ and $B \subseteq A_b^* \cup A_b^\omega$. If $w \in A_b^*$ and $i \geq 0$ is an integer, then w^i is the concatenation $ww \cdots w$ (i times) and w^ω is the infinite concatenation $ww \cdots w \cdots$. The \cdot operator can be omitted when the meaning is clear, as in $w\eta$.

By $w \sqsubseteq u$ and $w \sqsubset y$ we denote that w is a prefix of u and y , respectively. Further, let $\mathbf{pref}(y) = \{w : w \sqsubset y\}$ and $\mathbf{infix}(y) = \{w : \exists v(v \cdot w \sqsubset y)\}$ be the set of prefixes and infixes of y , respectively.

1.2 Preliminary definitions and results

In this section we define the classes of real numbers studied in the paper.

A real number α is called a *Liouville number* if it is irrational and for every positive integer k , there exist integers p_k and q_k with $q_k > 1$

such that

$$\left| \alpha - \frac{p_k}{q_k} \right| < \frac{1}{q_k^k}.$$

A real $\alpha \in [0, 1]$ is called *computable* if for some $b \geq 2$ its has a b -ary computable expansion $\alpha = 0.x_1x_2\dots$, that is, there is a computable function f_α such that $f_\alpha(n) = x_n$, for all $n \geq 1$. This condition is equivalent to the requirement that there is a computable sequence of rationals $(\frac{p_n}{q_n})_{n \in \mathbb{N}}$ such that $|\alpha - \frac{p_n}{q_n}| \leq \frac{1}{2^n}$, for all $n \in \mathbb{N}$. This shows that if α is computable, then its expansions in any base b are computable.

Originally, ω -words \mathbf{x} were called disjunctive because the syntactic monoid of the set $\{\mathbf{x}\}$ is disjunctive, that is, its syntactic congruence is the identity (see [JST83]). Equivalently, disjunctive ω -words are those which have every finite word as subword.¹ In fact, in a disjunctive ω -word every word appears infinitely many times.

Disjunctivity is also related to randomness: disjunctive ω -words are exactly the ω -words not contained in any null-set definable by finite automata [St76, St98]. For more properties of disjunctive sequences see [CPS97].

A real number $\alpha \in [0, 1]$ is *disjunctive* (or *rich*) in base b if its b -ary expansion is disjunctive. For example, Champernowne's number $0.0123456789101112\dots$ is a computable disjunctive in base 10 [Ch33]. No rational number is disjunctive in any base.

An *absolutely disjunctive* number (or *lexicon*) is a real which is disjunctive in every base. Every Martin-Löf random real is a lexicon, but the converse is false [Ca02].

In the sequel we denote by \mathcal{L} , \mathcal{C} and \mathcal{D} the set of all Liouville numbers, computable numbers and absolutely disjunctive numbers in $[0, 1]$, respectively.

1.3 Co-meagre and dense sets

It is useful to consider the unit interval $[0, 1]$ and the spaces of infinite sequences A_b^ω as metric spaces. Suitable metrics are the usual distance $|\alpha - \beta|$ in $[0, 1]$ and

$$\rho(\mathbf{x}, \mathbf{y}) = b^{-\inf\{i \in \mathbb{N} \mid i \geq 1, x_i \neq y_i\}},$$

¹In view of this latter property they are also called *rich* ω -words.

for infinite words $\mathbf{x} = x_1 \cdots x_i \cdots, \mathbf{y} = y_1 \cdots y_i \cdots$ with $x_i, y_i \in A_b$. With these metrics A_b^ω and $[0, 1]$ become complete metric spaces.

Let \mathcal{X} be a complete metric space. A set $M \subseteq \mathcal{X}$ is *nowhere dense* if its closure (smallest closed set containing M) does not contain a non-empty open subset. A set $M \subseteq \mathcal{X}$ is *meagre* (or of *first Baire category*) if it is a countable union of nowhere dense sets. A complement of a meagre set is called *co-meagre* (or *residual*).

The following closure property of co-meagre sets is well-known (see [Ku66]).

Fact 1 *In a complete metric space the family of co-meagre sets is closed under countable intersection.*

A set $M \subseteq \mathcal{X}$ is *dense* if $M \cap M' \neq \emptyset$ for every non-empty open set $M' \subseteq \mathcal{X}$. Note that in a complete metric space every co-meagre set is dense, but a dense set might be meagre, even countable.

The following relations hold for subsets $F \subseteq A_b^\omega$ and their counterparts in $[0, 1]$.

Lemma 2 ([St02]) *Let $F \subseteq A_b^\omega$ and $M_F = \{0.\mathbf{x} \mid \mathbf{x} \in F\} \subseteq [0, 1]$. Then*

1. *F is nowhere dense if and only if M_F is nowhere dense.*
2. *F is co-meagre if and only if M_F is co-meagre.*
3. *F is dense if and only if M_F is dense.*

Fact 3 ([Ox80]) 1. *The set of Liouville numbers \mathcal{L} is co-meagre.*

2. *The set of computable numbers \mathcal{C} is countable, meagre and dense.*

2 Disjunctive ω -words

As mentioned above disjunctive ω -words are infinite words $\mathbf{x} \in A_b^\omega$ having $\mathbf{infix}(\mathbf{x}) = A_b^*$. By

$$D_b = \{\mathbf{x} \mid \mathbf{x} \in A_b^\omega \wedge \mathbf{infix}(\mathbf{x}) = A_b^*\}$$

we denote the set of all disjunctive ω -words in A_b^ω . Then the set of all absolutely disjunctive numbers in $[0, 1]$ is

$$\mathcal{D} = \{\alpha \mid \alpha \in [0, 1] \wedge \forall b(b \geq 2 \rightarrow \exists \mathbf{x}(\mathbf{x} \in D_b \wedge \alpha = 0.\mathbf{x}))\}.$$

The set \mathcal{D} has the following topological property:

Lemma 4 ([CZ99, St02]) *The set \mathcal{D} is co-meagre in $[0, 1]$.*

Then from Fact 1 and Lemma 2 it follows that the set of absolutely disjunctive Liouville numbers is “topologically” large:

Corollary 5 *The intersection $\mathcal{L} \cap \mathcal{D}$ is co-meagre in $[0, 1]$.*

Corollary 5 gives only an existence proof, not a constructive one. Further more, since the set of computable reals \mathcal{C} is countable, it does not even show that $\mathcal{L} \cap \mathcal{D} \cap \mathcal{C} \neq \emptyset$.

To show the existence of computable absolutely disjunctive Liouville numbers we derive a representation of the b -ary counterparts $\{\mathbf{x} \in A_b^\omega \mid 0.\mathbf{x} \in \mathcal{D}\}$ of \mathcal{D} via computable languages. In Section 4 we then show how this description can be transformed into an algorithm computing an absolutely disjunctive Liouville number.

Theorem 6 ([St02]) *For every base b there effectively exists a computable language $W_b \subseteq A_b^*$ such that the ω -language $\{\mathbf{x} \in A_b^\omega \mid \text{the set } \mathbf{pref}(\mathbf{x}) \cap W_b \text{ is infinite}\}$ is the set of all b -ary expansions of absolutely disjunctive reals in $[0, 1]$.*

More explicitly, Theorem 6 ([St02, Theorem 21]) provides, for every base b , an increasing computable function $g : \mathbb{N} \rightarrow A_b^*$ such that $g(\mathbb{N}) = W_b$. This function g naturally induces a computable order on W_b .

Since \mathcal{D} is dense in $[0, 1]$, from Lemma 2.3 we deduce that that the ω -language $\{\mathbf{x} \in A_b^\omega \mid \text{the set } \mathbf{pref}(\mathbf{x}) \cap W_b \text{ is infinite}\}$ is dense in A_b^ω . This yields the following.

Corollary 7 *For every $u \in A_b^*$ there is a $v \in W_b$ such that $u \sqsubset v$.*

Proof. As the ω -language $\{\mathbf{x} \in A_b^\omega \mid \text{the set } \mathbf{pref}(\mathbf{x}) \cap W_b \text{ is infinite}\}$ is dense, hence every open subset of A_b^ω contains an \mathbf{x} such that $\mathbf{pref}(\mathbf{x}) \cap W_b$ is infinite.

Consider the open ω -language $u \cdot A_b^\omega$ (see e.g. [St02]). Then there is an \mathbf{x} for which $\mathbf{pref}(\mathbf{x}) \cap W_b$ is infinite. Consequently, there is a $v \in \mathbf{pref}(\mathbf{x}) \cap W_b$ such that $u \sqsubset v$. \square

3 Expansions of Liouville numbers

For our purposes it is useful to have the following property of b -ary expansions \mathbf{x} of reals which guarantees that $0.\mathbf{x}$ is a Liouville number. A similar criterion was sketched, without proof by Maillet in [Ma04].

Using finitely or infinitely many strings $w_i \in A_b^*$ and a function $f : \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}$ we construct b -ary expansions of real numbers in the following way.

Define $\Lambda_{j=0}^\infty w_j^{f(j)}$ as the concatenation of w_0 ($f(0)$ times), w_1 ($f(1)$ times), w_2 ($f(2)$ times)...

Lemma 8 ([CS17]) *Let $(w_i)_{i \in \mathbb{N}}$ be a family of non-empty strings $w_i \in A_b^*$, $f : \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}$, and $n_i = \sum_{j=0}^i f(j) \cdot |w_j|$. If*

$$\liminf_{i \rightarrow \infty} \frac{n_{i-1} + |w_i|}{n_{i-1} + f(i) \cdot |w_i|} = 0, \quad (1)$$

then $\mathbf{x} = \Lambda_{j=0}^\infty w_j^{f(j)}$ is the b -ary expansion of a rational or a Liouville number.

4 The Algorithm

The following algorithm computes the b -ary expansion $\mathbf{x} = \Lambda_{j=0}^\infty w_j^{f(j)}$ of an absolutely disjunctive Liouville number whose b -ary expansion starts with a given word $w_0 \in A_b^*$. It uses the computable injective ordering $g : \mathbb{N} \rightarrow W_b$ of the computable language W_b given by Theorem 6.

Algorithm Liouville-disjunctive

```

0   initialise  $w_0 = u_0 = v_0, \quad f(0) = 1$ 
1       for  $i = 1$  to  $\infty$  do
2            $v_i =$  first word in  $(W_b \cap u_{i-1} \cdot A_b^*) \setminus \{u_{i-1}\}$ 
3           calculate  $w_i$  from  $v_i = u_{i-1} \cdot w_i$ 
4           calculate  $f(i) = \min \left\{ k \mid \frac{|u_{i-1}| + |w_i|}{|u_{i-1}| + k \cdot |w_i|} < \frac{1}{i} \right\}$ 
5            $u_i = u_{i-1} \cdot w_i^{f(i)}$ 
6       endfor

```

The algorithm computes three families of words $(u_i)_{i \in \mathbb{N}}$, $(v_i)_{i \in \mathbb{N}}$, and $(w_i)_{i \in \mathbb{N}}$ and a function $f : \mathbb{N} \rightarrow \mathbb{N} \setminus \{0\}$. Note that at each step the set $(W_b \cap u_{i-1} \cdot A_b^*) \setminus \{u_{i-1}\}$ is effectively ordered according to g .

First, Step 2 implies $v_i \in W_b$ and together with Step 5, by induction, $u_{i-1} \sqsubset v_i \sqsubseteq u_i \sqsubset v_{i+1}$. From the Step 3 and $u_{i-1} \sqsubset v_i$ we have $|w_i| > 0$. Then, again using Step 5, by induction one verifies that $u_i = \Lambda_{j=0}^i w_j^{f(j)}$.

It remains to show that the algorithm will produce an infinite computable ω -word, that is, it never stops. To this end it suffices to show that the choice in Step 2 is always possible. From Corollary 7 we know that for every $u \in A_b^*$ there is a $v \in W_b$ such that $u \sqsubset v$. This makes it possible to choose the first element in W_b w.r.t. g which has u as a proper prefix.

Thus the algorithm computes two computable approximations of an ω -word $\mathbf{x} = \Lambda_{j=0}^\infty w_j^{f(j)}$ via the families of prefixes $(u_i)_{i \in \mathbb{N}}$ and $(v_i)_{i \in \mathbb{N}}$. From $v_i \in W_b$ we obtain $0.\mathbf{x} \in \mathcal{D}$ via Theorem 6, and, since $u_i = \Lambda_{j=0}^i w_j^{f(j)}$, Step 4 shows that the words u_i and w_i satisfy Eq. (1). Thus Lemma 8 verifies that $0.\mathbf{x}$ is also a Liouville number. The computability of \mathbf{x} follows directly from the algorithm.

Acknowledgement

The authors are grateful to Helmut Jürgensen for introducing them (long time ago) to Liouville numbers and disjunctive ω -words.

References

- [BHS15] V. BECHER, P. A. HEIBER, T. A. SLAMAN. A computable absolutely normal Liouville number, *Math. Comp.* **84** (2015) 296, 2939–2952.
- [Bu02] Y. BUGEAUD. Nombres de Liouville et nombres normaux, *C. R. Math. Acad. Sci. Paris* **335** (2002) 2, 117–120.
- [Ca02] C. S. CALUDE. *Information and Randomness: An Algorithmic Perspective*, Springer, Berlin, 2002. (2nd edition)
- [CPS97] C. S. CALUDE, L. PRIESE, L. STAIGER. Disjunctive Sequences: An Overview, *CDMTCS Research Report* 63, 1997, 40 pp.
- [CS17] C. S. CALUDE, L. STAIGER. Liouville, Computable, Borel normal and Martin-Löf random numbers, *Theory Comput. Syst.* (2017), to appear.
- [CZ99] C. S. CALUDE, T. ZAMFIRESCU. Most numbers obey no probability laws, *Publ. Math. Debrecen* **54** (1999) suppl., 619–623. Automata and formal languages, VIII (Salgótarján, 1996).
- [Cs59] J. W. S. CASSELS. On a problem of Steinhaus about normal numbers, *Colloq. Math.* **7** (1959), 95–101.
- [Ch33] D. G. CHAMPERNOWNE. The construction of decimals normal in the Scale of Ten. *J. London Math. Soc.* **S1-8**, 4 (1933), 254.
- [He96] P. HERTLING. Disjunctive ω -words and real numbers, *JUCS* **2** 7, (1996), 549–568.
- [JST83] H. JÜRGENSEN, H. J. SHYR, G. THIERRIN. Disjunctive ω -languages, *Elektron. Informationsverarb. Kybernet.* **19**, 6 (1983), 267–278.
- [JT88] H. JÜRGENSEN, G. THIERRIN. Some structural properties of ω -languages, In: *Proc. 13th Nat. School with Internat. Participation, Applications of Math. in Technology*, Sofia, 1988, 56–63.

- [Ku66] K. KURATOWSKI. *Topology. Vol. I*. Academic Press, New York-London and Państwowe Wydawnictwo Naukowe, Warsaw, 1966.
- [Ma04] E. MAILLET. Sur les nombres quasi-rationnels et les fractions arithmétiques ordinaires ou continues quasi-périodiques, *C. R. Acad. Sci., Paris* **138** (1904), 410–411.
- [Ox80] J. C. OXTOPY. *Measure and Category*, Springer-Verlag, New York-Berlin, 1980. (2nd edition)
- [Sc60] W. M. SCHMIDT. On normal numbers, *Pacific J. Math.* **10** (1960), 661–672.
- [St76] L. STAIGER. Reguläre Nullmengen, *Elektron. Informationsverarbeitung. Kybernetik* **12**, 6 (1976), 307–311.
- [St98] L. STAIGER. Rich ω -words and monadic second-order arithmetic, In: M. NIELSEN, W. THOMAS (eds.), *Computer science logic (Aarhus, 1997)*. Lecture Notes in Computer Science 1414, Springer-Verlag, Berlin, 1998, 478–490.
- [St02] L. STAIGER. How large is the set of disjunctive sequences? *J.UCS* **8**, 2 (2002) 348–362.
- [St02a] L. STAIGER. The Kolmogorov complexity of real numbers, *Theoret. Comput. Sci.* **284**, 2 (2002), 455–466.